

$SU(1, 1)$ group action and the corresponding Bloch equation

A. Okninski

Physics Division, Politechnika Swietokrzyska,
Al. 1000-lecia PP 7, 25-314 Kielce, Poland.

October 31, 2009

Abstract

Discrete time dynamics on the $SU(1, 1)$ group is studied. It is shown that a map acting in the dual space is the stroboscopic map for a $SU(1, 1)$ Bloch equation. Exact solution of the map is used to elucidate the corresponding dynamics. It is shown that dynamics of the $SU(1, 1)$ Bloch equation in the elliptic case bears close analogy to the $SU(2)$ Bloch dynamics.

1 Introduction

Discrete-time dynamical systems can be formulated in terms of group actions to exploit the group structure and get a better understanding of the corresponding dynamics. This approach was used to study discrete-time dynamics on some groups, see [1] and references therein. On the other hand, structure of Kleinian groups is naturally studied in the setting of discrete-time dynamical systems, revealing in this way connections with fractals [2, 3, 4]. For example, the Shimizu-Leutbecher map is a typical tool to study group structure [5, 6], see also [2, 3, 7].

Recently, we have investigated a possibility of relating group actions with stroboscopic maps of ordinary differential equations [1]. More exactly, we have studied the following dynamical system on a Lie group $\mathcal{G}$:

$$R_{N+1} = Q_N R_N Q_{N-1} R_{N-1} Q_{N-1}^{-1} R_N^{-1} Q_N^{-1}, \quad N = 1, 2, \dots, \quad (1)$$

where $Q_N, R_N \in \mathcal{G}$. A general solution of the map (1) has been constructed and it was demonstrated that for $\mathcal{G} = SU(2)$ and $Q_N \equiv Q$ Eq.(1) is a stroboscopic map of the Bloch equation [1]. The latter result is generalized in the present paper for the case $\mathcal{G} = SU(1, 1)$.

Let us note here that the $SU(1, 1)$ Bloch equation finds important applications in quantum optics. More exactly, the group of squeezings is generated by the Lie algebra $\mathfrak{su}(1, 1)$, the geometry of group manifold is that of

Minkowski spaces and time evolution is described by the $SU(1, 1)$ Bloch equation [8, 9, 10, 11].

The paper is organized as follows. In the next Section discrete time dynamics, defined and solved in [1] for arbitrary group $\mathcal{G}$, is studied in the case $\mathcal{G} = SU(1, 1)$. It is shown in Section 3 that the map (1), considered in the dual space, is the stroboscopic map for the $SU(1, 1)$ Bloch equation which is written in the elliptic, parabolic and hyperbolic cases. In Section 4 computations intended to elucidate dynamics of the $SU(1, 1)$ Bloch equation are presented for the elliptic case and analogy with the $SU(2)$ Bloch equation is stressed. The obtained results are summarized in the last Section.

2 Discrete-time dynamics on the $SU(1, 1)$ group

Let us recall that the map (1) admits an exact solution:

$$R_{2K} = S_{2K} S_{2K-1} \dots S_2 R_0 S_2^{-1} \dots S_{2K-1}^{-1} S_{2K}^{-1}, \quad (2a)$$

$$S_N = Q_{N-1} \dots Q_1 S_1 Q_{-1}^{-1} \dots Q_{N-3}^{-1}, \quad (2b)$$

where

$$S_N \stackrel{df}{=} R_N Q_{N-1} R_{N-1} Q_{N-2}, \quad (3)$$

and similar equations can be written for R_{2K+1} [1].

We shall consider a special case $Q_N \equiv Q$ in (1). In the case $\mathcal{G} = SU(1, 1)$ the following parameterization is used [12]:

$$R_N = \exp\left(i \frac{\chi_N}{2} \vec{\kappa} \cdot \vec{r}_N\right), \quad \vec{r}_N \cdot \vec{r}_N = \eta, \quad (4)$$

$$Q = \exp\left(i \frac{\alpha}{2} \vec{\kappa} \cdot \vec{q}\right), \quad \vec{q} \cdot \vec{q} = \eta, \quad (5)$$

where $i^2 = -1$, $\eta \in \{+1, 0, -1\}$ and is fixed, $\vec{\kappa} \stackrel{df}{=} [i\sigma^1, i\sigma^2, \sigma^3]$ where $\sigma^1, \sigma^2, \sigma^3$ are the Pauli matrices and scalar products are defined as

$$\vec{\kappa} \cdot \vec{x} \stackrel{df}{=} -\kappa^1 x^1 - \kappa^2 x^2 + \kappa^3 x^3, \quad (6a)$$

$$\vec{x} \cdot \vec{x} \stackrel{df}{=} -x^1 x^1 - x^2 x^2 + x^3 x^3, \quad (6b)$$

where $\vec{x} \stackrel{df}{=} [x^1, x^2, x^3]$. The three cases $\eta = +1, 0, -1$ are referred to as elliptic, parabolic and hyperbolic, respectively. For an exposition of theory of the $SU(1, 1)$ group the reader can consult [12, 13].

We obtain from (2) the following solution:

$$R_{2K} = Q^{2K} P^K R_0 P^{-K} Q^{-2K}, \quad (7)$$

where $P \stackrel{df}{=} Q^{-1} S_1 Q^{-1} = Q^{-1} R_1 Q R_0$, $K = 0, 1, 2, \dots$. We still have to impose initial condition R_0 while R_1 is computed as $R_1 = Q P R_0^{-1} Q^{-1}$.

Matrix P is parameterized in form

$$P = \exp\left(i \frac{\beta}{2} \vec{\kappa} \cdot \vec{p}\right), \quad \vec{p} \cdot \vec{p} = \eta, \quad (8)$$

and equation (7) can be written as

$$\vec{\kappa} \cdot \vec{r}_{2K} = \exp(iK\alpha \vec{\kappa} \cdot \vec{q}) \exp\left(iK\frac{\beta}{2} \vec{\kappa} \cdot \vec{p}\right) \vec{\kappa} \cdot \vec{r}_0 \exp\left(-iK\frac{\beta}{2} \vec{\kappa} \cdot \vec{p}\right) \exp(-iK\alpha \vec{\kappa} \cdot \vec{q}), \quad (9)$$

and, after introducing new quantities, $K\alpha = \theta$, $\lambda = \frac{\beta}{2\alpha}$, reads

$$\vec{\kappa} \cdot \vec{r}(\theta) = \exp(i\theta \vec{\kappa} \cdot \vec{q}) \exp(i\lambda\theta \vec{\kappa} \cdot \vec{p}) \vec{\kappa} \cdot \vec{r}_0 \exp(-i\lambda\theta \vec{\kappa} \cdot \vec{p}) \exp(-i\theta \vec{\kappa} \cdot \vec{q}), \quad (10)$$

where $\vec{r}(\theta) \stackrel{df}{=} \vec{r}_{2K}$.

2.1 Elliptic case

To obtain $\vec{r}(\theta)$ from Eq.(10) we shall need to compute $\vec{t}(\gamma)$ given by:

$$\vec{\kappa} \cdot \vec{t}(\gamma) = S \vec{\kappa} \cdot \vec{t} S^{-1} = \exp\left(i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}\right) \vec{\kappa} \cdot \vec{t} \exp\left(-i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}\right), \quad (11)$$

in the case $\eta = 1$. The exponential map is simplified as

$$\exp\left(i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}\right) = \cos\left(\frac{\gamma}{2}\right) \mathbf{1} + i \sin\left(\frac{\gamma}{2}\right) \vec{\kappa} \cdot \vec{s}, \quad \vec{s} \cdot \vec{s} = 1. \quad (12)$$

Now, due to properties of the Pauli matrices we obtain from (11):

$$\vec{t}(\gamma) = \cos(\gamma) \vec{t} - \sin(\gamma) \vec{t} \times \vec{s} + (1 - \cos(\gamma)) (\vec{t} \cdot \vec{s}) \vec{s}, \quad (13)$$

where

$$\vec{x} \times \vec{y} \stackrel{df}{=} [-x_2y_3 + x_3y_2, -x_3y_1 + x_1y_3, x_1y_2 - x_2y_1]. \quad (14)$$

Using twice the equation (13) in (10) we get:

$$\vec{r}(\theta) = \cos(2\theta) \vec{t}(\theta) - \sin(2\theta) \vec{t}(\theta) \times \vec{q} + (1 - \cos(2\theta)) (\vec{q} \cdot \vec{t}(\theta)) \vec{q}, \quad (15a)$$

$$\vec{t}(\theta) = \cos(2\lambda\theta) \vec{r}_0 - \sin(2\lambda\theta) \vec{r}_0 \times \vec{p} + (1 - \cos(2\lambda\theta)) (\vec{p} \cdot \vec{r}_0) \vec{p}. \quad (15b)$$

For growing θ the vector $\vec{r}(\theta)$ evolves on manifold $\vec{r}(\theta) \cdot \vec{r}(\theta) = 1$, i.e. on two-sheeted hyperboloid.

2.2 Parabolic case

In the case $\eta = 0$ the exponential map reduces to

$$\exp\left(i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}\right) = \mathbf{1} + i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}, \quad \vec{s} \cdot \vec{s} = 0. \quad (16)$$

Now, due to properties of the Pauli matrices we compute from (11):

$$\vec{t}(\gamma) = \vec{t} - \gamma \vec{t} \times \vec{s} + (\vec{t} \cdot \vec{s}) \vec{s}. \quad (17)$$

Using twice the equation (17) in (10) we obtain:

$$\vec{r}(\theta) = \vec{t}(\theta) - 2\theta \vec{t}(\theta) \times \vec{q} + (\vec{q} \cdot \vec{t}(\theta)) \vec{q}, \quad (18a)$$

$$\vec{t}(\theta) = \vec{r}_0 - 2\lambda\theta \vec{r}_0 \times \vec{p} + (\vec{p} \cdot \vec{r}_0) \vec{p}. \quad (18b)$$

The vector $\vec{r}(\theta)$ evolves on manifold $\vec{r}(\theta) \cdot \vec{r}(\theta) = 0$, i.e. on the Minkowski cone.

2.3 Hyperbolic case

In the case $\eta = -1$ we have

$$\exp(i\frac{\gamma}{2} \vec{\kappa} \cdot \vec{s}) = \cosh(\frac{\gamma}{2}) \mathbf{1} + i \sinh(\frac{\gamma}{2}) \vec{\kappa} \cdot \vec{s}, \quad \vec{s} \cdot \vec{s} = -1. \quad (19)$$

Due to properties of the Pauli matrices we obtain from (11):

$$\vec{t}(\gamma) = \cosh(\gamma) \vec{t} - \sinh(\gamma) \vec{t} \times \vec{s} + (\cosh(\beta) - 1) (\vec{t} \cdot \vec{s}) \vec{s}. \quad (20)$$

Using twice the equation (20) in (10) we get:

$$\vec{r}(\theta) = \cosh(2\theta) \vec{t}(\theta) - \sinh(2\theta) \vec{t}(\theta) \times \vec{q} + (\cosh(2\theta) - 1) (\vec{q} \cdot \vec{t}(\theta)) \vec{q}, \quad (21a)$$

$$\vec{t}(\theta) = \cosh(2\lambda\theta) \vec{r}_0 - \sinh(2\lambda\theta) \vec{r}_0 \times \vec{p} + (\cosh(2\lambda\theta) - 1) (\vec{p} \cdot \vec{r}_0) \vec{p}. \quad (21b)$$

The vector $\vec{r}(\theta)$ evolves on manifold $\vec{r}(\theta) \cdot \vec{r}(\theta) = -1$, i.e. on one-sheeted hyperboloid.

2.4 Symmetry and restrictions of dynamics

Dynamical system (1) for $Q_N \equiv Q$ has continuous symmetry:

$$R_N \rightarrow Q^\kappa R_N Q^{-\kappa}, \quad \forall \kappa \in \mathbb{R}. \quad (22)$$

It can be thus expected that dynamics of the quantity $\vec{r}(\theta) \cdot \vec{q}$ should decouple from other degrees of freedom in (1) [1]. Indeed, it follows from (15) that

$$\vec{r}(\theta) \cdot \vec{q} = \vec{t}(\theta) \cdot \vec{q}. \quad (23)$$

Since in the elliptic or hyperbolic case $\vec{t} \cdot \vec{t} = \vec{q} \cdot \vec{q} = \pm 1$ it follows from the Schwartz inequality for the Minkowski metric that $(\vec{t} \cdot \vec{q})^2 \geq (\vec{t} \cdot \vec{t})(\vec{q} \cdot \vec{q}) =$
1. Now, for given $\vec{p}$, $\vec{q}$ and $\vec{r}_0$ on the upper sheet of the hyperboloid in the elliptic case we have

$$1 \leq A_1 \leq \vec{t}(\theta) \cdot \vec{q} \leq A_2, \quad (24)$$

The constants $A_{1,2}$ depending on the parameters $\vec{p}$, $\vec{q}$ and the initial condition $\vec{r}_0$ can be computed from (15b) by elementary means

$$A_{1,2} = c \mp \sqrt{b^2 + (a - c)^2}, \quad (25)$$

where

$$a = \vec{r}_0 \cdot \vec{q}, \quad b = (\vec{r}_0 \times \vec{p}) \cdot \vec{q}, \quad c = (\vec{p} \cdot \vec{r}_0) (\vec{p} \cdot \vec{q}). \quad (26)$$

It thus follows that the motion on the hyperboloid is bounded by two planes: $A_1 \leq \vec{r}(\theta) \cdot \vec{q} \leq A_2$. In the parabolic case we obtain the following simple condition

$$\vec{t} \cdot \vec{q} = a - 2\lambda\theta b + c, \quad (27)$$

i.e. equation of a straight line.

3 The Bloch equation

The map (1), $Q_N \equiv Q$, is the stroboscopic map of a differential equation which is conveniently deduced from the form (10). Since α and β are arbitrary we shall treat θ as a continuous variable. Differentiating Eq.(10) with respect to θ and using (10) we get

$$\frac{d \vec{\kappa} \cdot \vec{r}(\theta)}{d\theta} = i [\vec{\kappa} \cdot \vec{u}(\theta), \vec{\kappa} \cdot \vec{r}(\theta)], \quad (28)$$

where $[A, B] \stackrel{df}{=} AB - BA$ and

$$\vec{u}(\theta) = \vec{q} + \lambda \vec{p}(\theta), \quad (\lambda = \beta/2\alpha) \quad (29a)$$

$$\vec{\kappa} \cdot \vec{p}(\theta) = \exp(i\theta \vec{\kappa} \cdot \vec{q}) \vec{\kappa} \cdot \vec{p} \exp(-i\theta \vec{\kappa} \cdot \vec{q}). \quad (29b)$$

It follows that the sequence $\vec{r}_0, \vec{r}_2, \vec{r}_4, \dots$, generated by $R_0, R_2, R_4, \dots$, cf. Eq.(7), interpolates flow of Eq.(28). It turns out that, θ interpreted as time, Eq.(28) is the $SU(1,1)$ Bloch equation, cf.[10].

Equations (28), (29) can be written in explicit form.

3.1 Elliptic case

Using Eqs. (11), (13) we get from (29b)

$$\vec{p}(\theta) = \cos(\gamma) \vec{p} - \sin(\gamma) \vec{p} \times \vec{q} + (1 - \cos(\gamma)) (\vec{p} \cdot \vec{q}) \vec{q}, \quad (30)$$

and using properties of the Pauli matrices we obtain from (28) the Bloch equation in the elliptic case:

$$\frac{d \vec{r}}{d\theta} = -2 \vec{r}(\theta) \times \vec{u}(\theta), \quad (31)$$

with $\vec{u}(\theta)$ and $\vec{p}(\theta)$ given by (29a) and (30), respectively.

3.2 Parabolic case

Applying Eqs. (11), (17) to (29b)

$$\vec{p}(\theta) = \vec{p} - \gamma \vec{p} \times \vec{q} + (\vec{p} \cdot \vec{q}) \vec{q}, \quad (32)$$

and using properties of the Pauli matrices we obtain from (28) the Bloch equation in the parabolic case:

$$\frac{d\vec{r}}{d\theta} = -2\vec{r}(\theta) \times \vec{u}(\theta), \quad (33)$$

with $\vec{u}(\theta)$ and $\vec{p}(\theta)$ given by (29a) and (32), respectively.

3.3 Hyperbolic case

Using Eqs. (11), (20) we obtain from (29b)

$$\vec{p}(\theta) = \cosh(\gamma) \vec{p} - \sinh(\gamma) \vec{p} \times \vec{q} + (\cosh(\gamma) - 1) (\vec{p} \cdot \vec{q}) \vec{q}, \quad (34)$$

and using properties of the Pauli matrices we obtain from (28) the Bloch equation in the hyperbolic case:

$$\frac{d\vec{r}}{d\theta} = -2\vec{r}(\theta) \times \vec{u}(\theta), \quad (35)$$

with $\vec{u}(\theta)$ and $\vec{p}(\theta)$ given by (29a) and (34), respectively.

4 Computational results

We have performed several computations for the Bloch equation (31) and the discrete-time dynamical system (1), parameterized as in Eqs. (4), (5), $Q_N \equiv Q$, $\eta = 1$. Exact solutions of the map (1) as well as of the Bloch equation (31) in the elliptic case are given by (7) and (15), respectively.

The solution (15), of discrete-time dynamical system (1) with Q, P given by (5), (8) has been plotted in Fig. 1 for $\vec{q} = [0, 0, 1]$, $\vec{p} = [1, 0, \sqrt{2}]$, $\lambda = \beta/(2\alpha) = 3$ and the initial vector $\vec{r}_0 = [\frac{1}{2}, \frac{1}{2}, \sqrt{\frac{3}{2}}]$ on the upper sheet of the hyperboloid. The whole trajectory has three-fold symmetry with respect to the $\vec{q}$ axis. Circles indicate parallels $A_{1,2}$ given by (25) confining the dynamics.

In Fig. 2 dynamics of vectors $\vec{r}_N$ obtained from (9) has been plotted for $\alpha = 5$, $\beta = 20$ ($\lambda = \beta/(2\alpha) = 2$) and other parameters unchanged. We thus obtain thirty six points marked with dots. The solution (15) has been also plotted. The closed curve has two-fold symmetry with respect to the $\vec{q}$ axis.




Fig. 1. Exact solution of the Bloch equation (31), $\lambda = 3$.




Fig. 2. Exact solution of the Bloch equation (31) (thin line) and discrete-time dynamical system (1) (dots), $\alpha = 5$, $\beta = 20$, $\lambda = 2$.

In Fig. 3 initial stage of dynamics of vectors $\vec{r}_N$ has been plotted for $\lambda = \beta / (2\alpha) = 1.025$, dot marking the initial vector $\vec{r}_N$.




Fig. 3. Exact solution of the Bloch equation (31), $\lambda = 1.025$.

5 Summary and discussion

We have introduced in [1] a class of discrete-time invertible maps (1) on an arbitrary group $\mathcal{G}$ for which the exact solution (2) of this map has been found. Maps of form (1), parameterized on a Lie group, generate points in the dual (parameter) space which sample a trajectory in this space arbitrarily densely. This curve can be generated forward as well as backward from a given initial condition. This suggests that the group action (1) may correspond to a flow of a differential equation. We have demonstrated in the present paper that for $\mathcal{G} = SU(1,1)$ the map (1), $Q_N \equiv Q$, considered as dynamical system in the dual space, is a stroboscopic map of a $SU(1,1)$ Bloch equation. It should be noted that the $SU(1,1)$ Bloch equation (28) is formally analogous to the $SU(2)$ Bloch equation, cf. Eq. (4.21) in [1].

Exact solutions of the map constructed in the present paper, (15), (18), (21), lead to a better understanding of the corresponding Bloch equations (31), (33), (35). More exactly, symmetries and restrictions of dynamics have been found explicitly. It is interesting that dynamics of the $SU(1,1)$ Bloch equation in the elliptic case bears close analogy to dynamics of the $SU(2)$ Bloch equation, compare Figs. 1, 2, 3 from the present paper with analogous figures in [1].

References

- [1] A. Okninski, Acta Phys. Polon. B **40** (2009) 1605-1616; arXiv:0804.2128v3 [math-ph].
- [2] R. Brooks, J.P. Matelski, in *Riemann Surfaces and Related Topics: Proceedings of the 1978 Stony Brook Conference*, edited by I.Kra, B. Maskit, *Ann. Math. Studies* **97** (Princeton Univ. Press, Princeton, N.J., 1981), p. 65.
- [3] F.W. Gehring, G.J. Martin, Bull. Am. Math. Soc. **21** (1989) 57-63.
- [4] D. Mumford, C. Series, D. Wright, *Indra's pearls: The vision of Felix Klein*. Cambridge University Press, Cambridge, 2002.
- [5] H. Shimizu, Ann. of Math. **77** (1963) 33-71.
- [6] A. Leutbecher, Math. Zeit. **100** (1967) 183-200.
- [7] A.F. Beardon, *The Geometry of Discrete Groups*. Springer-Verlag, 1983.
- [8] G. Dattoli, A. Dipace, A. Torre, Phys. Rev. A **33** (1986) 4387-4389.
- [9] P.K. Aravind, J. Opt. Soc. Am. B **5** (1988) 1545-1553.
- [10] B.E. King, *Quantum State Engineering and Information Processing with Trapped Ions*, PhD Thesis, Department of Physics, University of Colorado, 1999 (<http://jilawww.colorado.edu/pubs/thesis/king/>).
- [11] R.R. Puri, *Mathematical Methods of Quantum Optics*. Springer, Berlin 2001.
- [12] G. Chiribella, G.M. D'Ariano, P. Perinotti, Laser Physics **16** (2006) 1572-1581; arXiv:0610142v1 [quant-ph].
- [13] A. Wawrzyńczyk, *Group representations and special functions*. D. Reidel Publishing Co., Dordrecht/Boston/Lancaster, 1984.